A Case of the Therapeutic Effect of *Nangan-jeon* on Refractory Functional Dyspepsia Caused by “Liver-Kidney Deficiency Cold”

Chang-gue Son

Liver and Immunology Research Center, Oriental Medical College in Daejeon University

A Case of the Therapeutic Effect of *Nangan-jeon* on Refractory Functional Dyspepsia Caused by “Liver-Kidney Deficiency Cold”

Chang-gue Son

Liver and Immunology Research Center, Oriental Medical College in Daejeon University

ABSTRACT

Objective: This study explored the clinical usefulness of *Nangan-jeon* decoction to treat refractory functional dyspepsia.

Methods: A 61-year-old female had been suffering from severe functional dyspepsia for about 40 years, but the dyspeptic symptoms rapidly improved after taking *Nangan-jeon* decoction. The clinical outcome was assessed by the numerical rating scale (NRS) and the self-reporting method.

Result: After there was no improvement from various herbal treatments in a department of a Korean college hospital, the patient moved to the author's clinic. She had been diagnosed with the pattern identification of “liver-kidney deficiency cold” and prescribed with *Nangan-jeon* decoction. This herbal drug rapidly improved her dyspeptic symptoms, from NRS 10 to NRS 2 (after 3 weeks) and NRS 1 (after 7 weeks).

Conclusion: This case report is the first to present the potential of *Nangan-jeon* decoction to improve refractory functional dyspepsia.

Key words: traditional Korean medicine, functional dyspepsia, *Nangan-jeon*

I . Introduction

Functional dyspepsia (FD) is one of most common disorders in gastrointestinal (GI) system, as the

estimated prevalence of 10% to 30% worldwide and 8 to 37 % in Korea¹⁻³. It refers to the recurrent or persistent dyspeptic symptoms more than 6 months without explainable organic causes⁴. Accordingly, the diagnosis of FD is made by the self-report-based feature of symptoms called ‘Rome IV Criteria for functional criteria’⁵. FD is known as a multi-causal disorder, which infection of *Helicobacter pylori*, visceral hypersensitivity, low-grade mucosal inflammation, and/or delayed gastric emptying are involved in its pathophysiology⁶.

· 투고일: 2021.05.12, 심사일: 2021.06.21, 게재확정일: 2021.06.21
· Corresponding author: Chang-gue Son Liver and Immunology Research Center, Daejeon Oriental Hospital of Daejeon University, 75 Daedeok Daero 176 beon-gil, Seo-gu, Daejeon, Republic of Korea
TEL: +82-42-470-9134
FAX: +82-42-470-9005
E-mail: ckson@dju.ac.kr

FD patients have a poor quality of life (QoL) more than those with other GI functional disorders including gastroesophageal reflux or irritable bowel syndrome⁷. In general, proton pump inhibitors (PPI), selective serotonin reuptake inhibitors (SSRIs), tricyclic antidepressants (TCA) or prokinetic drugs are used for treatments of FD⁸. FD has the highly complex pathologies, which induce these conventional therapies to have limitations such as lower response and frequent relapse after cessation⁹. Accordingly, many clinicians and patients are looking for herbal remedies in an aspect of multi-targeted medicine¹⁰. For example, *Banhasasim-tang* (半夏瀉心湯) and *Yukgunja-tang* (六君子湯), two popular herbal prescriptions had shown the clinical efficacy on FD patients^{11,12}.

On the other hand, a quarter of patients has a refractory FD characterized by a long duration, severe symptoms, and poor QoL in spite of medical treatment¹³. These refractory FD patients are particularly problematic, and many alternative approaches including electroacupuncture have been tried to manage them¹⁴. In addition, a patient-customized herbal therapy is recommended for these patients.

This study reports a severe case of refractory FD who showed a rapid response to an *Nangan-jeon* (暖肝煎) decoction. *Nangan-jeon* a formula for 'cold symptoms of the liver- and kidney-Yin' (肝腎虛寒), then no study exist for the ant-FD therapeutic effect yet.

II. Report of the case

1. Characteristics of patient and medical history

A 61-year-old woman has been suffering from persisting postprandial fullness, early satiation after meal and sometimes epigastric pain since her

teenage. The symptoms have used to restrict her food life and to worsen daily life frequently. She had been diagnosed as functional dyspepsia long before visiting author's clinic, based on non-specific finding from gastroscopy multiple times. The patient frequently had visited Western clinics and took various conventional drugs (prokinetics, PPI or SSRIs) when she felt the bad condition. However, her symptoms had been repeated the a little better and worsen again. From April 2020, the patient felt the worse dyspepsia-related symptoms after hard exercise and heavy sweating over.

The patient was generally susceptible to cold and psychological stress, and no change of body weight with around 19.0 of body-mass index (height 157 cm, body weight 47 kg). She had no specific family history, but had a diagnosis of Raynaud's disease in her 20s. The patient thus has tried to physical exercise intentionally, and has neither used alcohol, smoking and drugs, nor had food-allergic reaction.

This case study was approved by the institutional review board (IRB) at Daejeon Korean Medicine Hospital (IRB number: DJDSKH-21-E-14-1).

2. Diagnosis, treatments, and course of symptom

Her abdominal discomfort including postprandial fullness, gastric satiation and especially severe epigastric pain became worse from one year, while these symptoms were affected by eating meals and worsened under cold circumstance. Her complaints weren't better even various treatments from both Western and Korean medicine clinics, and thus she was transferred to author's clinic on January 2021. As expected, her symptoms met the Rome IV Criteria for functional criteria¹⁴ (Table 1). On September 2020, the patient started to be treated

in other department of same hospital for about four months, but not much improved even six different herbal decoctions, *Hyangsapyungwi-san* (香砂平胃散), *Hyangsayukgunja-tang* (香砂六君子湯), *Banhahubak-tang* (半夏厚朴湯), *Pyeongjingeonbi-tang* (平陳健脾胃湯), *Gyeji-tang* (桂枝湯) and *Yikgeobohyul-tang* (益氣補血湯), respectively.

Author examined her history and characteristics of clinical symptoms, while she had a mild white-color coat at tongue and rapid-weak pulsation. The patient fell into *Soumin* of Sasang constitutional classification using QSCCII. Author thought the cold-derived dyspepsia based on the pattern identifications of “*liver-kidney deficiency cold* (肝腎虛寒)”. Author tried to test the response to *Ojeok-san* (五積散), and then confirmed the quick positive effect. The self-reporting NRS score (0 means no symptom while 10 doses the worst severity¹⁵) was quickly

decreased from NRS 10 to NRS 7 in 2 days. This positive response gave author the convincing diagnosis of “*Liver-Kidney Deficiency Cold*” which decided to prescribe *Nangan-jeon* (Table 2). This herbal drug rapidly improved her overall dyspeptic symptoms, into NRS 5 (after 1 week) and NRS 2 (after 3 weeks) followed by normal level of digestive function (NRS 1) after 7-week treatment, respectively (Fig. 1). In addition, her cold-sensitivity also decreased by NRS 5 on 4-week and gradually became better in length of taking *Nangan-jeon* (Fig. 1). In fact, acupuncture (CV12, HT7, LI4, SP4, SP6, ST36), moxibustion (CV4, KI1) and *Aconitum* pharmacopuncture (CV4 and BL23) were also given to her, as twice weekly for one month and followed by once per week later, respectively. As expected, her mild white-color coat at tongue was changed into a bright-color and the rapid-weak pulsation also a little improved.

Table 1. Rome IV Criteria for Functional Criteria

Criteria fulfilled for the last 3 months with symptom onset at least 6 months before diagnosis. No evidence of structural disease (including at upper endoscopy) that is likely to explain the symptoms and, must include one or more of:

1. Bothersome postprandial fullness
2. Bothersome early satiation
3. Bothersome epigastric pain
4. Bothersome epigastric burning

Table 2. Composition of *Nangan-jeon*

Scientific name	Herbal name	Chinese name	Dose*
<i>Lycium chinense</i> Miller	<i>Lycii Fructus</i>	枸杞子	24.0 g
<i>Angelica gigas</i> Nakai	<i>Angelicae Gigantis Radix</i>	當歸	16.0 g
<i>Poria cocos</i> Wolf	<i>Poria Sclerotium</i>	白茯苓	16.0 g
<i>Foeniculum vulgare</i> Miller	<i>Foeniculi Fructus</i>	小茴香	16.0 g
<i>Lindera strichnifolia</i> Fernandez-Villar	<i>Linderae Radix</i>	烏藥	16.0 g
<i>Cinnamomum cassia</i> Presl	<i>Cinnamomi Cortex</i>	肉桂	8.0 g
<i>Aquilaria agallocha</i> Roxburgh	<i>Aquilariae Agallochae Lignum Resinatum</i>	沈香	4.0 g

* The indicated dose is weight of each herb for an adult during one day

Fig. 1. Summary of clinical outcome and treatment courses.

III. Discussion and Conclusion

FD is known to show the epidemiologic features of a 1.4 to 2.7-fold female predominance along with a lower QoL than men¹⁶. Author also found a higher prevalence of FD in females by approximately 2-folds (14.2% of male versus 26.9% in female)¹⁷. In addition, females were known to poorly respond in FD by *H. pylori* eradication, one of main treatments for FD¹⁸. A clinical data using 1,341 Chinese FD patients presented 24.4% of the refractory FD, and a tendency of female-predominance (male 22.9% versus 25.4%)¹². In general, we call refractory FD if symptoms are unresponsive to at least two medical treatments, such as acid suppressors, proton pump inhibitors, or prokinetics, for six months¹⁹.

The patient of this case report showed a typical refractory FD pattern because both previous conventional and herbal drugs didn't work efficiently. In aspect of the subtypes of FD, this patient showed the overlapping feature of postprandial distress syndrome (PDS) and epigastric pain syndrome (EPS). In a Korean study, the proportion of PDS and EPS

were 53.6% and 31.8%, while 14.6% of FD belonged to the PDS plus EPS²⁰. One clinical study revealed the preference of the EPS+PDS subtype in patients with refractory FD than in patients with non-refractory FD¹².

This study assessed the improvement of symptoms using self-reported NRS score, then her most of signs including postprandial fullness and notable epigastric pain rapidly decreased. The previously prescribed herbal decoctions, *Hyangsapyungwi-san*, *Hyangsayukgunja-tang*, *Banhahubak-tang*, and *Pyeongjingeonbi-tang*, are generally used for FD, while some have been proven for their anti-FD efficacy from clinical studies^{21,22}. This patient was however refractory to these decoctions. In Korean medicine, FD can be generally classified into 6 different pattern identifications such as 'spleen and stomach deficiency and cold', 'liver-stomach disharmony', 'spleen deficiency with qi stagnation', 'tangled cold and heat', 'dampness and heat in the spleen and stomach systems' or 'food retention disorder'²³. Then, 'liver-kidney deficiency cold', the patient's pattern identification was not included.

In fact, the choose of the *Nangan-jeon* was made by the rapid response to *Ojeok-san*. *Ojeok-san* usually treats the pain-related disorders, but it has also the pharmacological action on the 'cold'-related dyspepsia. In previous department, other two decoctions (*Gyeji-tang* and *Yikgeebohyul-tang*) targeting each the 'cold'-related symptoms and 'deficiency of qi and blood' weren't effective unlikely *Nangan-jeon*. *Nangan-jeon* first appeared in *Gyeongakjeonseo* (景岳全書), as a formula for 'cold symptoms of the liver- and kidney-Yin'²⁴. This formula showed the pharmaceutical properties to improve the female-fertility, under situations of especially 'liver-kidney cold type'^{25,26}, which supported the rationale to choose *Nangan-jeon* this woman patient. As 'cold'-focused treatments were added using electronic moxibustion (CV4, KI1) and *Aconitum* pharmacopuncture (CV4 and BL23), as well as *Soumin*-related acupuncture (CV12, HT7, LI4, SP4, SP6, ST36), respectively. *Aconitum* was reported to exhibit a warming effect on the abdominal region such as CV12 and CV8²⁷, accordingly these treatments together might contribute to the improvement of this case.

To date, many approaches including electroacupuncture and psychiatric management using a tricyclic antidepressant have been tried to solve the refractory FD, but many limitations are still existing^{28,29}. In summary, even this is only a case, *Nangan-jeon*-based therapeutics could be a fine choice for refractory FD. Of course, further studies are required to verify the therapeutic effects of *Nangan-jeon* and the proper applicable type of refractory FD.

References

1. Stanghellini V, Chan FK, Hasler WL, Malagelada JR, Suzuki H, Tack J, et al. Gastrointestinal disorders. *Gastroenterology* 2016;150(5):1380-92.
2. Mahadeva S, Goh KL. Epidemiology of functional dyspepsia: a global perspective. *World J Gastroenterol* 2006;12(17):2661-6.
3. Jung HK, Keum BR, Jo YJ, Jee SR, Rhee PL, Kang YW, et al. Diagnosis of functional dyspepsia: a systematic review. *Korean J Gastroenterol* 2010;55(5):296-307.
4. Stanghellini V, Chan FKL, Hasler WL, Malagelada JR, Suzuki H, Tack J, et al. Gastrointestinal disorders. *Gastroenterol* 2016;150(6):1380-92.
5. Suzuki H. The Application of the Rome IV Criteria to Functional Esophagogastrroduodenal Disorders in Asia. *J Neurogastroenterol Motil* 2017;23(3):325-33.
6. Talley N, Walker M. Emerging evidence that irritable bowel syndrome & functional dyspepsia are microbial diseases. *Indian J Med Res* 2019; 149(4):437-40.
7. El-Serag HB, Talley NJ. Health-related quality of life in functional dyspepsia. *Aliment Pharmacol Ther* 2003;18(4):387-93.
8. Tack J, Camilleri M. New developments in the treatment of gastroparesis and functional dyspepsia. *Curr Opin Pharmacol* 2018;43:111-7.
9. Lacy BE, Talley NJ, Locke GR, Bouras EP, DiBaise JK, El-Serag HB, et al. Review article: current treatment options and management of functional dyspepsia. *Aliment Pharmacol Ther* 2012;36(1):3-15.
10. Chiarioni G, Pesce M, Fantin A, and Sarnelli G. Complementary and alternative treatment in functional dyspepsia. *United Eur Gastroenterol J* 2018;6(1):5-12.
11. Kim SK, Joung JY, Ahn YC, Jung IC, Son CG. Beneficial Potential of Banha-Sasim-Tang for

- Stress-Sensitive Functional Dyspepsia via Modulation of Ghrelin: A Randomized Controlled Trial. *Front Pharmacol* 2021;12:636752.
12. Tominaga K, Sakata Y, Kusunoki H, Odaka T, Sakurai K, Kawamura O, et al. Rikkunshito simultaneously improves dyspepsia correlated with anxiety in patients with functional dyspepsia: A randomized clinical trial (the DREAM study). *Neurogastroenterol Motil* 2018;30(7):e13319.
13. Jiang SM, Jia L, Lei XG, Xu M, Wang SB, Liu J, et al. Incidence and psychological-behavioral characteristics of refractory functional dyspepsia: a large, multi-center, prospective investigation from China. *World J Gastroenterol* 2015;21(6):1932-7.
14. Zheng H, Xu J, Sun X, Zeng F, Li Y, Wu X, et al. Electroacupuncture for patients with refractory functional dyspepsia: A randomized controlled trial. *Neurogastroenterol Motil* 2018;30(7):e13316.
15. Hartrick CT, Kovan JP, Shapiro S. The numeric rating scale for clinical pain measurement: a ratio measure? *Pain Pract* 2003;3(4):310-6.
16. Kim YS, Kim N. Functional Dyspepsia: A Narrative Review with a Focus on Sex-Gender Differences. *J Neurogastroenterol Motil* 2020;26(3):322-34.
17. Kim YJ, Ahn YC, Son CG. Sasang constitution affects the prevalence of functional dyspepsia. *BMC Complement Altern Med* 2015;15:150.
18. Kim SE, Kim N, Park SM, Kim WH, Baik GH, Jo Y, et al. Female gender is a poor predictive factor of functional dyspepsia resolution after *Helicobacter pylori* eradication: A Prospective, Multi-center Korean Trial. *Korean J Gastroenterol* 2018;72(6):286-94.
19. Hamilton J, Guthrie E, Creed F, Thompson D, Tomenson B, Bennett R, et al. A randomized controlled trial of psychotherapy in patients with chronic functional dyspepsia. *Gastroenterology* 2000;119(3):661-9.
20. Kim SE, Park HK, Kim N, Joo YE, Baik GH, Shin JE, et al. Prevalence and risk factors of functional dyspepsia: a nationwide multicenter prospective study in Korea. Prospective, Multi-center Korean Trial. *J Clin Gastroenterol* 2014;48(2):e12-8.
21. Kim JB, Shin JW, Kang JY, Son CG, Kang W, Lee HW, et al. A traditional herbal formula, Hyangsa-Pyeongwi san (HPS), improves quality of life (QoL) of the patient with functional dyspepsia (FD): randomized double-blinded controlled trial. *J Ethnopharmacol* 2014;151(1):279-86.
22. Lv L, Wang FY, Ma XX, Li ZH, Huang SP, Shi ZH, et al. Efficacy and safety of Xiangsha Liujunzi granules for functional dyspepsia: A multi-center randomized double-blind placebo-controlled clinical study. *World J Gastroenterol* 2017;23(30):5589-601.
23. Ha NY. Development of Standard Tool for Pattern Identification of Functional Dyspepsia and Analysis of its Clinical Characterization. *Graduate School of Korean Medicine, Kyung Hee University, PhD's thesis* 2020.
24. He J, Huang M, Zhang Q, Liao W, Wang L, Zhang Y. Treatment of unstable angina pectoris with modified nuan gan jian - a report of 33 cases. *J Tradit Chin Med* 2004;24(4):263-5.
25. Lee SH, Lee YH. Case Reports of Treatment of Nangan-jeon for Female Infertility. *J Naturopathy* 2013;2(1):83-8.
26. Lee SH. Influences of Gagamanganjeon and Grain Syrup on the Infertile Women with Symptoms

- of Liver-Kidney-Cool Types. *J Naturopathy* 2014;4(1):88-98.
27. Yang JM, Li G, Wang M, Jin YX, Zheng FJ, Sun Y, et al. The Effects of Natural Chinese Medicine Aconite Root, Dried Ginger Rhizome, and Coptis on Rectal and Skin Temperatures at Acupuncture Points. *Evid Based Complement Alternat Med* 2017;2017:7250340.
28. Chung VC, Wong CH, Wu IX, Ching JY, Cheung WK, Yip BH, et al. Electroacupuncture plus on-demand gastrocaine for refractory functional dyspepsia: Pragmatic randomized trial. *J Gastroenterol Hepatol* 2019;34(12):2077-85.
29. Cheong PK, Ford AC, Cheung CKY, Ching JYL, Chan Y, Sung JJY, et al. Low-dose imipramine for refractory functional dyspepsia: a randomized, double-blind, placebo-controlled trial. *Lancet Gastroenterol Hepatol* 2018;3(12):837-44.